Colorado MCH Trend Analysis 2012

Ashley Juhl, MSPH and Sue Ricketts, Ph.D March 8, 2013



What is the state trend analysis?

A spreadsheet that shows

- Short term and long term trends in maternal and child health indicators
- Healthy People 2020 and Colorado 2020 targets
- Progress
- Problems


Everything you want to know about 49 indicators

- Preconception/prenatal
- Infants
- Children
- Teens
- Oral health




State of Colorado Trend Analysis, 2012 Progress Toward Healthy People 2020 Targets or Colorado 2020 Targets

| T) | Frogress roward healthy reopie 2020 rargets or Colorado 2020 rargets | | | | | | | | | | |
|---|--|-------------------|------------------|-------|-------|-------|-------|--------------------|-----------------------|------------------------------------|-----------------------------------|
| Maternal and Child Health Indicator | | Healthy People | Year | | | | | | Short Term Trend: | Long Term Trend: Percent Change | 2010 Position Relative to 2020 |
| | Source | 2020 Target | 2000 | 2006 | 2007 | 2008 | 2009 | 2010 | 2006 to 2010 | 2000 to 2010 | Target |
| Preconception/Prenatal | | | | | | | | | | | |
| % mothers who smoked 3 months before pregnancy | BC | 14.6% | NA | NA | 11.2% | 11.1% | 10.7% | 10.4% | Improved ^a | NA NA | Meets target |
| % mothers gained appropriate weight during pregnancy b | BC | 38% ^c | NA | NA | 31.2% | 31.6% | 32.3% | 33.1% ^d | | NA | Close to target |
| % mothers overweight or obese before pregnancy ^b | ВС | 38% ^c | NA | NA | 35.0% | 34.8% | 35.9% | 43.2% ^d | NA | NA | Close to target |
| % mothers received adequate prenatal care | BC | 77.6% | 68.4% | 74.0% | 67.0% | 67.9% | 68.5% | 69.1% | Worse | +1.0% | Close to target |
| % births unintended | PRAMS | 33% ^c | 40% | 40% | 36% | 37% | 40% | 36% | No real change | -10% | Close to target |
| % women 18-44 mental health not good in past 30 days | BRFSS | 40% ^c | 48% | 41% | 45% | 42% | 43% | 47% | Worse | -2% | Close to target |
| % women 18-44 with any physical activity in last 30 days | BRFSS | 90% ^c | 78% | 82% | 83% | 82% | 84% | 82% | No real change | +5% | Close to target |
| % women and men 18-44 using effective birth control | BRFSS | 80% ^c | 68% ^e | 65% | NA | NA | 67% | 68% | No real change | 0% ⁹ | Close to target |
| % mothers who smoked during last 3 months of pregnancy | BC | 1.4% | NA | NA | 7.3% | 7.2% | 6.7% | 6.5% | Improved ^a | NA NA | Far from target |
| % mothers drinking alcohol last 3 months pregnancy | PRAMS | 3 1.7% | 9% | 11% | 11% | 11% | 11% | 12% | No real change | +29% | Far from target |
| % mothers with excessive weight gain during pregnancy | BC | NA. | NA | NA | 44.8% | 47.2% | 49.4% | 47.7% ^d | NA | NA | NA |
| % women 18-44 eating 5+ fruits and vegetables daily | BRFSS | S NA | 23% | NA | 30% | NA | 27% | NA | NA | NA | NA |
| Unemployment rate (%), women with children under 6 | ACS | NA. | NA | NA | 7% | 5% | 9% | 10% | Worse | NA | NA |
| Infants | 11 == 7 | / | | | | | | | | | |
| % mothers initiating breastfeeding | PRAMS | 88% ^{c*} | 86% | 90% | 90% | 90% | 92% | 93% | No real change | +9% | Meets target |
| % mothers breastfeeding at 6 months | CHS | 60.6% | 51% ^f | 54% | 56% | 60% | 61% | 61% | Improved | +20% ^h | Meets target |
| % very low birth weight births in Level IIIs | BC | 83.7% | 74.8% | 87.5% | 87.1% | 88.3% | 87.8% | 89.2% | Improved | +19.3% | Meets target |
| Infant mortality rate | BC/DC | 6.0 | 6.1 | 5.7 | 6.1 | 6.2 | 6.3 | 5.9 | Worse | -3.3% | Meets target |
| Postneonatal mortality rate | DC | 2.0 | 1.9 | 1.3 | 1.9 | 1.8 | 2.1 | 1.6 | Worse | -15,8% | Meets target |
| Newborn hearing screening (%) | EHDI | 98% ^{c*} | 90.7% | 97.6% | 97.2% | 97.8% | 97.3% | 97.3% | Worse | +7.3% | Close to target |
| % births low birth weight | BC | 7.8% | 8.5% | 9.0% | 9.0% | 8.9% | 8.8% | 8.8% | Improved | +3.5% | Close to target |
| Neonatal mortality rate | DC | 4.1 | 4.3 | 4.4 | 4.2 | 4.4 | 4.2 | 4.3 | Improved | 0.0% | Close to target |
| Black/white infant mortality ratio | DC | NA. | NA | 2.3 | 2.3 | 2.5 | 2.9 | 2.4 | Worse | NA | NA |



Preview

Distance to 2020 Target for 35 Indicators





Healthy People 2010 vs. 2020 Targets

| Indicator | HP 2010 Target | HP 2020 Target | | | | |
|--------------------------|-------------------|-------------------|--|--|--|--|
| Easier to Meet | | | | | | |
| Adequate prenatal care | 90.0% | 77.6% | | | | |
| Infant mortality rate | 4.5 | 6.0 | | | | |
| Low birth weight | 5.0% | 7.8% | | | | |
| Motor vehicle death rate | 9.2 | 12.4 | | | | |
| Neonatal death rate | 2.9 | 4.1 | | | | |
| Postneonatal death rate | 1.2 | 2.0 | | | | |
| Suicide death rate | 5.0 | 10.2 | | | | |
| Harder to Meet | | | | | | |
| Breastfeeding at 6 mos. | 50.0% | 60.6% | | | | |
| Child abuse rate | 10.3 | 8.5 | | | | |



2010 Data Relative to Targets


| Indicator | Distance from HP 2010 | Distance from HP 2020 | | | | | | |
|--------------------------|--------------------------|--------------------------|--|--|--|--|--|--|
| Easier to Meet | | | | | | | | |
| Adequate prenatal care | At some distance | Close to target | | | | | | |
| Infant mortality rate | At some distance | Meets target | | | | | | |
| Low birth weight | At some distance | Close to target | | | | | | |
| Motor vehicle death rate | At some distance | Meets target | | | | | | |
| Neonatal death rate | At some distance | Close to target | | | | | | |
| Postneonatal death rate | At some distance | Meets target | | | | | | |
| Suicide death rate | Far from target | Close to target | | | | | | |
| Harder to Meet | | | | | | | | |
| Breastfeeding at 6 mos. | Meets target | Meets target | | | | | | |
| Child abuse rate | Meets target | Close to target | | | | | | |





Preconception/Prenatal

Distance to 2020 Target for 10 Indicators



3 indicators do not have a 2020 target



Preconception/Prenatal (continued)

Short Term Trend

- Right direction
 - Smoked 3 months before pregnancy
 - Smoked during last 3 months of pregnancy
- Wrong direction
 - Adequate prenatal care
 - Poor mental health
 - Unemployment for women with children under 6



Preconception/Prenatal (continued)

- No real change
 - Unintended births
 - Physical activity
 - Using effective birth control
 - Drinking during last 3 months of pregnancy
- Need more data to determine trends
 - Appropriate weight gain during pregnancy
 - Overweight/obese before pregnancy
 - Excessive weight gain during pregnancy
 - Fruit and vegetable consumption




Preconception/Prenatal (continued)

- Long Term Trend
 - Unintended births improved
 - Drinking during the last three months of pregnancy worsened



Infants

Distance to 2020 Target for 8 Indicators



One indicator does not have a 2020 target



Infants (continued)

Short Term Trend

- Right direction
 - Breastfeeding at 6 months
 - Very low birth weight births in level IIIs
 - Low birth weight births
 - Neonatal mortality rate
- No real change
 - Breastfeeding initiation

Infants (continued)


- Wrong direction
 - Infant mortality rate
 - Postneonatal mortality rate
 - Newborn hearing screening
 - Black/White infant mortality ratio
- Long Term Trend
 - Breastfeeding at 6 months improved
 - Very low birth weight births in Level IIIs improved





Children

Distance to 2020 Target for 9 Indicators



4 indicators do not have a 2020 target



Children (continued)

Short Term Trend

- Right direction
 - Injury hospitalization rate 0-14
 - Motor vehicle death rate 0-14
 - Overweight/obese children
 - Child abuse rate
 - Health insurance coverage
 - Sugar-sweetened beverage consumption
 - Fruit and vegetable consumption
 - CHP+ enrollees
 - Medicaid enrollees
 - ▶ Child death rate I-I4



Children (continued)

- Mixed
 - ▶ Parents asked to fill out a questionnaire on child
- Wrong direction
 - ▶ Children 0-18 at or below 100% federal poverty level
- Need more data to determine trend
 - ▶ Children I-I4 with a medical home



Children (continued)

Long Term Trend


- Injury hospitalization rate 0-14 improved
- Motor vehicle death rate 0-14 improved
- ▶ Child death rate I-I4 improved
- Children 0-18 at or below 100% federal poverty level worsened





Teens

Distance to 2020 Target for 8 Indicators



4 indicators do not have a 2020 target



Teens (continued)

Short Term Trend

- Right direction
 - Injury hospitalization rate
 - Motor vehicle death rate
 - Physical activity
 - Using effective birth control
 - Teen fertility rate
- Wrong direction
 - Graduation rate
 - ▶ Teen suicide death rate
 - Chlamydia rate



Teens (continued)

- Need more data to determine trend
 - Participation in extracurricular activities
 - Sugar-sweetened beverage consumption
 - Eating breakfast every day
 - School enjoyment

Long Term Trend

- Injury hospitalization rate improved
- Motor vehicle death rate improved
- ▶ Teen fertility rate improved
- Chlamydia rate worsened



Oral Health

- ▶ 2 indicators
 - No 2020 targets
- Short Term Trend
 - Right direction
 - Medicaid dental served
 - No real change
 - Children who went to a dentist by age I

In Conclusion

Summary

- Colorado is meeting many 2020 targets already
 - But remember many are easier to meet now
- The majority of indicators showed improvement over the past five years (2006-2010)
- Nine indicators showed improvement since 2000; three were worse than in 2000 (where measurable)



In Conclusion (continued)

What Needs to Happen Next

- Maintain indicators meeting the targets
- Work to improve those
 - Not meeting the targets
 - Going in the wrong direction





Data Sources

- Colorado Health and Environmental Data (CHED)
 - http://www.chd.dphe.state.co.us/default.aspx
- Colorado Health Information Dataset (CoHID)
 - http://www.chd.dphe.state.co.us/cohid/Default.aspx
- Children with Special Health Care Needs
 - http://www.colorado.gov/cs/Satellite/CDPHE-PSD/CBON/1251617590646



Any Questions?

